

Evidence for a role of the oxytocin system, indexed by genetic variation in *CD38*, in the social bonding effects of expressed gratitude

Sara B. Algoe^{1,*} and Baldwin M. Way^{2,*}

¹Department of Psychology, University of North Carolina at Chapel Hill, Chapel Hill, NC 27599, USA, and ²Department of Psychology and the Ohio State Institute for Behavioral Medicine Research, The Ohio State University, Columbus, Ohio 43210

Oxytocin is thought to play a central role in promoting close social bonds via influence on social interactions. The current investigation targeted interactions involving expressed gratitude between members of romantic relationships because recent evidence suggests gratitude and its expression provides behavioral and psychological ‘glue’ to bind individuals closer together. Specifically, we took a genetic approach to test the hypothesis that social interactions involving expressed gratitude would be associated with variation in a gene, *CD38*, which has been shown to affect oxytocin secretion. A polymorphism (rs6449182) that affects *CD38* expression was significantly associated with global relationship satisfaction, perceived partner responsiveness and positive emotions (particularly love) after lab-based interactions, observed behavioral expression of gratitude toward a romantic partner in the lab, and frequency of expressed gratitude in daily life. A separate polymorphism in *CD38* (rs3796863) previously associated with plasma oxytocin levels and social engagement was also associated with perceived responsiveness in the benefactor after an expression of gratitude. The combined influence of the two polymorphisms was associated with a broad range of gratitude-related behaviors and feelings. The consistent pattern of findings suggests that the oxytocin system is associated with solidifying the glue that binds adults into meaningful and important relationships.

Keywords: close relationships; gratitude; emotion expression; *CD38*; oxytocin

INTRODUCTION

For centuries, scholars have pondered the nature of the ties that bind people together. In the 1950s, psychologists Harlow (1958) and Bowlby (1958) emphasized the necessity of maternal emotional warmth and comfort for healthy child development. More recently, theorists have posited that the biological, behavioral and psychological systems that co-evolved in support of parent–child attachments have come to support close adult relationships as well (e.g. Hazan and Diamond, 2000; Diamond, 2004). One approach to illuminating the mechanisms through which people form and maintain these important relationships has been to shine the empirical spotlight on common social interactions that occur within them. Recent evidence strongly implicates the emotion and expression of gratitude in the adult bonding process (Algoe, 2012; Algoe *et al.*, 2013). In this study, we test whether oxytocin may support the behavior and acute psychological response surrounding the expression of gratitude in real time. We do this by using a genetic indicator of oxytocin secretion, variation in the *CD38* gene, and by creating a meaningful psychological context in which oxytocin is likely to reveal its influence: live interactions between members of ongoing romantic relationships.

THE BONDING EFFECTS OF GRATITUDE

Recent work regarding the emotion of gratitude supports the theory that this momentary emotional response to someone else’s kind

gesture has evolved to connect people more closely to others who would make high-quality relationship partners (see review in Algoe, 2012). Earlier models of gratitude’s role in social life focused on economic factors like costs to the benefit-provider (i.e. benefactor), value to the recipient and repayment of the benefactor (e.g. Trivers, 1971; McCullough *et al.*, 2001). However, the most recent evidence clarifies that, rather than merely facilitating interpersonal accounting of resources, gratitude works through a particular relational currency that is associated with intimacy, perceived partner responsiveness (Algoe *et al.*, 2008). Perceived partner responsiveness underlies relational intimacy and is associated with feeling understood, accepted and cared for (Reis *et al.*, 2004); it is an important situational trigger for feeling gratitude upon receipt of a benefit (Algoe *et al.*, 2008; Algoe and Stanton, 2012). In turn, the grateful individual is likely to demonstrate responsiveness for the benefactor’s own needs and preferences in the future (e.g. Kubacka *et al.*, 2011). In short, the positive emotion of gratitude provides fuel for upward spirals of mutual responsiveness between dyad members, thereby promoting the quality of the relationship over time.

In light of these strong connections between experienced emotion and relational outcomes, more recent work has begun to focus in on how one person’s experienced gratitude might translate to relational growth for each dyad member. One candidate behavior that follows from emotion theory is the expression of gratitude; indeed, in situations that would cause gratitude, acknowledging a benefit is the most frequently reported motive of the benefit recipient (Algoe and Haidt, 2009). Recent studies provide suggestive evidence that, for the grateful individual, expressing gratitude is necessary to reap the relational benefits (e.g. Lambert *et al.*, 2010; Algoe and Stanton, 2012). For example, participants who were randomly assigned to ‘go the extra mile’ to express gratitude to a friend over the course of 3 weeks reported greater communal relationship strength with that friend by the end of the study, even compared with a condition in which people were assigned to focus on thinking about things they appreciate about the friend over the 3-week period (Lambert *et al.*, 2010).

Received 20 May 2013; Revised 17 September 2013; Accepted 28 December 2013

Advance Access publication 5 January 2014

*These authors contributed equally to this work.

This work was supported by a grant from the National Institute of Mental Health (MH59615) and a grant from the National Center for Advancing Translational Sciences (8KL2TR000112-05). The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health. The authors would like to thank Ms Sara DeMaria for her excellent technical assistance, as well as the exceptional team of 30 research assistants who helped with the Carolina Couples Study, lead by Jenny Bridgers (full list available on first author’s website). Barbara Fredrickson, B. Keith Payne and Crystal Schiller provided insightful feedback on an earlier manuscript draft.

Correspondence should be addressed to Sara B. Algeo, Department of Psychology, University of North Carolina at Chapel Hill, CB #3270, Chapel Hill, NC 27599, USA. E-mail: algoe@unc.edu

Beyond the grateful person, though, because of gratitude's 'other-praising' nature (Algoe and Haidt, 2009), expressed gratitude may provide psychological rewards to the target of the expression (i.e. the original benefactor) that act as a hook to keep the benefactor engaged in and satisfied with the relationship with the grateful individual. Critically, this is most likely to be true if the expresser is perceived to be responsive when thanking the original benefactor (Algoe *et al.*, 2013). Specifically, a recent study used a naturalistic interaction in which one romantic partner expressed gratitude to the other in the lab; for targets of the expressions who perceived their romantic partner as being particularly responsive—that is, understanding, validating and caring (Reis *et al.*, 2004)—when expressing gratitude to them, the target's satisfaction with the relationship improved over the subsequent 6 months (Algoe *et al.*, 2013). In fact, effects of perceived partner responsiveness after the partner expressed gratitude to the target in the lab even held when statistically controlling for perceptions of the partner's responsiveness after participating in two different types of common couple interactions that have been documented to be important for relationship well-being (i.e. providing support for positive or negative events; Gable *et al.*, 2006). These findings highlight the unique and powerful role that expressions of gratitude can play in social bonding, beyond the other types of behaviors that may foster relationship quality. Identifying potential biological contributors to these effects stands to open doors for improved understanding of social bonding processes.

CLOSE RELATIONSHIPS AND OXYTOCIN

Since the seminal discovery that oxytocin facilitates monogamous bonds in rodents (Williams *et al.*, 1994), a rapidly growing body of research in humans has sought to determine if oxytocin is involved in human social connections, broadly defined. Accordingly, oxytocin has been associated with everything from parenting behavior (e.g. Feldman *et al.*, 2012) to behavior toward a new acquaintance (e.g. Kosfeld *et al.*, 2005). However, for the question of close social bonds, it is notable that the adult romantic relationship context is unique in its qualitative experience as well as its impact on health and well-being (e.g. Berscheid, 1999; Hazan and Diamond, 2000). Because the connection between oxytocin and psychosocial processes appears to be highly dependent on the behavioral and relational context (see also Bartz *et al.*, 2011), it is important to study psychosocial correlates of oxytocin directly in the romantic relationship context.

Within this context, the oxytocin system has been hypothesized to both facilitate bonding processes and change in response to social interactions (see Carter and Porges, 2013). In this work, we focus on the former type of research question, regarding the facilitation of bonding (for examples of the latter, see Grewen *et al.*, 2005; Holt-Lunstad *et al.*, 2008; Smith *et al.*, 2013). The few studies that have addressed whether or how oxytocin might facilitate bonding processes in the context of romantic relationships do show concurrent associations between the oxytocin system and a handful of relationship-relevant behaviors, as observed in well-defined laboratory interactions between couple-members. Specifically, intranasal administration of oxytocin led to an increase in positive communication during a conflict discussion (Ditzen *et al.*, 2009); similarly, higher levels of plasma oxytocin have been associated with greater positive behaviors during a conversation involving one person's expression of worry or concern (Gouin *et al.*, 2010), as well as while discussing a shared positive experience (Schneiderman *et al.*, 2012). Integrating across these diverse methodologies, these studies suggest that higher synaptic levels of oxytocin promote relationship building behaviors. Finally, variation in the oxytocin receptor gene (*OXTR*), which theoretically alters the impact of oxytocin signaling, was associated with greater use of affiliative cues

during a partner's expression of worry (Kogan *et al.*, 2011). These studies suggest that in general, the oxytocin system is involved in relationship-relevant processes. However, the mixture of methods used to measure or manipulate the oxytocin system combined with theory and evidence from relationship science suggesting that the various relationship behaviors investigated across these studies play different or unique roles in the promotion, maintenance and prevention-of-deterioration of the relationship (see Gable and Reis, 2001; Gable *et al.*, 2012; McNulty and Fincham, 2012), mean it is too early to draw broad conclusions about oxytocin's role in facilitating processes central to the promotion of close relationships.¹ In short, there is much more to learn.

As reviewed, within the adult close relationship context, there has been surprisingly little examination of early assertions (Uvnas-Moberg, 1996, 1998; Carter, 1998) regarding oxytocin's role in facilitating human social bonds—assertions which have been recently recapitulated (Carter and Porges, 2013). We believe a focus on social interactions involving expressed gratitude provides a prime opportunity to test associations between the oxytocin system and social processes involved in promoting adult human pair bonds because of the unique role of expressed gratitude in this process (Algoe *et al.*, 2013). Moreover, we go beyond the behavior of expressed gratitude to specify some of the acute psychological impacts of such interactions that should be particularly instrumental in the bonding process, namely positive emotions and perceived partner responsiveness.

GENETIC ASSOCIATIONS WITH RELATIONSHIP PROCESSES

As mentioned earlier, prior research suggests that genetic association methodology is a viable method for gaining clues to potential neurochemical underpinnings of social bonding related processes in romantic couples (Kogan *et al.*, 2011). We adopt a similar approach here. However, the cellular mechanism explaining how the previously discussed *OXTR* genetic variant could impact cellular signaling has not yet been identified, and a recent meta-analysis reported no association with a broad spectrum of socially relevant behaviors (Bakermans-Kranenburg and van IJzendoorn, 2014). Instead, we studied functional variation in a different gene, *CD38*, which is a key regulator of oxytocin release. Mice with deletion of the *CD38* gene exhibit marked reductions of oxytocin within the cerebrospinal fluid and the plasma (Jin *et al.*, 2007), indicating that *CD38* is necessary for oxytocin secretion. Unable to release oxytocin, these mice exhibit profound deficits in basic social processes such as social recognition and memory (Jin *et al.*, 2007).

In humans, *CD38* appears to have a role in oxytocin signaling and social processes as well. Levels of *CD38* gene expression in peripheral cells are positively correlated with plasma levels of oxytocin (Kiss *et al.*, 2011), suggesting that *CD38* facilitates oxytocin release. Commonly occurring variation in *CD38* (rs3796863) has also been associated with plasma oxytocin levels (Feldman *et al.*, 2012). Therefore, *CD38* is an important regulator of oxytocin signaling.

Consistent with a role in regulating oxytocin release, *CD38* is also associated with social processes. Accordingly, levels of peripheral *CD38* gene expression are related to clinician-observed social skills in autistic patients (Kiss *et al.*, 2011; Riebold *et al.*, 2011). The previously mentioned single nucleotide polymorphism (SNP) (rs3796863) has also

¹ To add complexity, another body of work tests concurrent associations between plasma oxytocin and self-reported assessment of global relationship evaluation rather than specific social behavior or its immediate impact in a live and meaningful context. Presumably, the quality of specific interpersonal interactions adds up to comprise the overall evaluation of the relationship. Yet in this line of work, researchers sometimes find negative associations between oxytocin and global relationship evaluation (e.g. Taylor, Saphire-Bernstein, & Seeman, 2010; Smith *et al.*, 2013). Though not the primary focus of the current investigation, we include a measure of global relationship satisfaction.

been associated with either low functioning autism (Lerer *et al.*, 2010) or high functioning autism (Munesue *et al.*, 2010). In the non-clinical context, a study of parenting behavior showed that rs3796863 was associated with reduced parental touch of infants (Feldman *et al.*, 2012), suggesting that *CD38* affects social engagement. Thus, there is an emerging literature tying rs3796863 to social processes. However, the molecular pathway by which this SNP could affect *CD38* expression or function has not been identified.

A different polymorphism in *CD38* that is located at the opposite end (5') of the *CD38* gene from rs3796863 has a more clearly delineated molecular pathway by which it can affect *CD38* expression. This SNP, rs6449182, is located in intron 1, which is a regulatory control region of the *CD38* gene (Ferrero *et al.*, 1999) and differentially affects transcription factor binding (Saborit-Villarroya *et al.*, 2011). Accordingly, this polymorphism has been associated with differences in relative levels of *CD38* messenger RNA (Jamroziak *et al.*, 2009), protein (Jamroziak *et al.*, 2009) and enzymatic activity (Polzonetti *et al.*, 2012). Thus, rs6449182 has functional effects on expression of *CD38* and, by extension, extracellular oxytocin levels in the brain and plasma. Although the mechanism by which rs6449182 affects *CD38* function is better delineated than for rs3796863, it has been less studied in a psychological context. Therefore in this study, we examine the association of rs6449182 as well as rs3796863 with gratitude and its social bonding effects. Because these two SNPs are located in different portions of the *CD38* gene, and genotypes at each locus are potentially independent (meaning that knowledge of one genotype does not necessarily predict the other), we also studied their combined effect by summing genotypes to create a putative index of *CD38* gene expression (see Chapman *et al.*, 2003 for this approach).

THIS STUDY

The current investigation focuses on whether variation in *CD38* is associated with social interactions involving the expression of gratitude. Evidence for the potential relational benefits from expressing gratitude (e.g. Lambert *et al.*, 2010) as well as receiving an expression of gratitude (Grant and Gino, 2010; Algoe *et al.*, 2013) suggests that the oxytocin system may influence both social roles: the expresser of gratitude and the recipient of gratitude expression. Moreover, we specify the relational and affective consequences of such interactions that are most closely related with high-quality, close, social connections: perceived partner responsiveness and positive emotions (see Fredrickson, 1998 for theory and Kok *et al.*, 2013 for the role of positive emotions in building social resources, over time). In addition, work that differentiates among positive emotions in momentary social function (e.g. Algoe and Haidt, 2009; Shiota *et al.*, 2011) suggests that it is worthwhile to further explore whether a specific active emotional ingredient in the romantic bonding context—love—is relevant (and see Gonzaga *et al.*, 2006 for links between romantic love and oxytocin). Thus, after receiving an expression of gratitude and after expressing gratitude, we examined perceptions of partner responsiveness as well as positive and negative emotions experienced; we also examined grateful behavior itself, observed in the lab and as gathered from daily self-reports.

METHOD

Participants

Each member of 77 heterosexual romantic couples participated in a study on 'Everyday experiences and feelings of people in romantic relationships' (aka the Carolina Couples Study; see additional information in Algoe *et al.*, 2013). Couples were recruited from the region surrounding Chapel Hill, North Carolina, and must have been romantically involved for a minimum of 6 months. The current sample consists of 128 individuals who provided saliva for genotyping (83.1% of

sample; 63 women, 65 men across 69 heterosexual couples; aged 18–57, $M = 28.5$, $s.d. = 8.65$). The majority of participants self-identified as White/Caucasian (76.2%), and the remaining participants were Black/African-American (11.9%), East or South Asian (4%), and multi-racial or of an unidentified racial category (7.9%); 3.2% identified as Hispanic. Couples had been together for about 4 years ($M = 4.2$ years), and were either dating exclusively (55.8%) or committed for life (i.e. married, engaged to be married or living as married; 44.2%).

Overview of design and procedure

The larger Carolina Couples Study was an observational investigation consisting of two visits to the lab, 2 weeks apart, with brief nightly questionnaires completed for each of 14 nights between lab visits. The current investigation focuses on one pair of 5 min interactions between couple-members at the second lab visit, in which participants were each given the opportunity to express gratitude to the partner, as well as nightly self-reported gratitude expression.

Expressed gratitude behavioral task

Each participant was asked to choose something specific, big or small, that the partner did for him or her, for which the participant felt grateful (see Algoe *et al.*, 2013). After noting the event, one participant took up to 5 min to thank the partner, after which they both self-reported their reactions to the interaction via private online questionnaires. The interaction and self-report were repeated with the second partner. Whether the male or female participant thanked the partner first was counterbalanced. Note that each participant in the study was part of two interactions: one in which he or she *expressed gratitude*, and one in which he or she *received an expression of gratitude*. Responses to each are important to our investigation because they reflect giving and receiving of positively valenced, other-focused attention, respectively.

Measures

At study entry, each participant reported on global relationship satisfaction using a well-validated seven-item measure (Hendrick, 1988; $\alpha = 0.82$).

After each interaction, participants rated their agreement with several items on scales ranging from 'not at all true/never true' (0) to 'very true/true all of the time' (6). First, each participant responded to 24 items to measure emotional response. To reduce demand effects, we included a broad range of items and information that any given situation may produce its own set of emotions. The 11 positive emotion terms (e.g. peaceful, loving, amused, proud) were averaged into a positive emotions composite ($\alpha = 0.90$, 0.86 after receiving an expression of gratitude and providing an expression, respectively). Note that the composite scores of the 13 negative emotion terms had extremely low ratings after each interaction ($M < 0.31$), consistent with the nature of the interactions, and we did not consider them further. Next, each participant responded to 10 items to measure the core components of perceived responsiveness (Gable *et al.*, 2006), which include understanding, validating and caring (10-item composite $\alpha = 0.93$, 0.95 after receiving an expression of gratitude and providing an expression, respectively). Data analysis included three measures both after expressing gratitude and after receiving an expression of gratitude: composite scores of perceived partner responsiveness and positive emotions, and, to probe specificity regarding theory, one targeted positive emotion term, 'loving'.

Grateful behavior was indexed by a participant's degree of praising of the partner's actions within the expression of gratitude (for theory, see Algoe and Haidt, 2009), and encompassed both verbal and non-verbal behaviors to guard against counting insincere praise. The behavior was reliably observed by four judges who rated the degree of

praising on a scale ranging from 1 (no or little praise) to 5 (excellent degree of benefactor praiseworthiness; $\alpha = 0.78$), and participants varied in their spontaneous use of praising behavior when expressing gratitude, with the average of judges' scores ranging from 1.25 to 5 ($M = 3.26$; $s.d. = 0.73$).

Finally, one question from the 14 nightly reports completed prior to the lab visit measured grateful behavior in a different way: each night, participants responded to the item, 'I thanked my partner for something he/she did that I appreciated' with a yes or no (coded as 1 and 0 for data analysis). This second measure provides the opportunity to test convergent and ecological validity of our hypothesis.

Genotyping

DNA was extracted from saliva collected using Oragene kits (DNA Genotek; www.dnagenotek.com) according to the manufacturer's instructions. Both SNPs were genotyped using Taqman SNP Genotyping Assays (rs6449182: C__1216863_10; rs3796863: C__1216944_10) from Applied Biosystems using an ABI 7300 Sequence Detection System. All samples were successfully genotyped on two separate occasions with complete concordance. Haploview (version 4.2; Barrett *et al.*, 2005) was used to calculate Hardy-Weinberg Equilibrium values using the exact test (Wigginton *et al.*, 2005) as well as the normalized measure of allelic association (D' ; Lewontin, 1964) and coefficient of determination (r^2 ; Hill and Robertson, 1968). The rs3796863 SNP was coded in a dominant manner [CC genotype = 0; A allele carriers (AC and AA) = 1] as in prior work (Feldman *et al.*, 2012, 2013; Sauer *et al.*, 2012). The CC genotype, which is present in approximately 40% of the Caucasian and East Asian populations, has been associated with lower CD38 expression (Lerer *et al.*, 2010) and lower plasma oxytocin levels (Feldman *et al.*, 2012). With respect to rs6449182, the G allele is generally associated with greater CD38 expression, which is reflected in higher levels of CD38 mRNA, protein, and enzymatic activity (Jamroziak *et al.*, 2009; Polzonetti *et al.*, 2012; but see Riebold *et al.*, 2011). This evidence on CD38 expression suggests that the G allele functions in an additive, not dominant manner, with the GG genotype (~5–10% of the Caucasian population) associated with the greatest expression. Therefore, the rs6449182 polymorphism was coded in an additive manner (CC = 0, CG = 1; GG = 2). To create a cumulative index of putative CD38 expression, the two SNPs were summed according to their effects on gene expression, as coded earlier.

RESULTS

Analysis plan

Table 1 contains descriptive statistics for all dependent measures. Dependent measures violate assumptions of independence required for typical ANOVA (analysis of variance) because each member of the couple provided reports on the same outcomes. Therefore, we use multilevel modeling (using HLM; Raudenbush *et al.*, 1996) for all analyses. Dependent measures from lab-based interactions were tested with two-level models (i.e. person nested within couple), and daily grateful behavior was tested with a three-level model (i.e. day within person within couple). (See Supplementary Table S1 containing correlations among dependent measures as well as intraclass correlations between dyads on each variable.) The distribution of genotypes for the two SNPs did not deviate from Hardy-Weinberg equilibrium ($P_s > 0.19$; for rs6449182- CC $n = 83$, CG $n = 37$, GG $n = 8$; for rs3796863, CC $n = 62$, A carriers $n = 66$). Models were run with rs6449182, rs3796863, or their combination as a continuous predictor variable for that person's own behavior or response to an interaction. None of the effects reported below was moderated by participant sex, nor were conclusions altered when controlling for ethnicity.

Table 1 Descriptive statistics for all dependent measures

	Mean	s.d.
Global relationship satisfaction	6.07	0.48
Receiving a gratitude expression		
Perceived partner responsiveness	5.06	0.75
Positive emotions	3.78	1.08
Loving	4.92	0.97
Expressing gratitude		
Perceived partner responsiveness	5.14	0.79
Positive emotions	4.11	0.91
Loving	5.00	1.01
Praising behavior	3.25	0.54
Daily expressed gratitude	0.79	0.12

Mean and s.d. reflect intercept and standard deviation of r , respectively, in unconditional HLM models where the listed variable was the dependent variable. For daily expressed gratitude, the intercept accounts for Bernoulli distribution of the outcome (0 or 1) and standard error (not deviation) is reported.

Similarity of rs6449182 and rs3796863

As expected due to their distinct locations in the CD38 gene, there was a relatively low degree of association between the two SNPs (in the primary ethnic group: $D' = 0.50$, 95% CI [0.09, 0.79]; overall: $D' = 0.45$, 95% CI [0.08, 0.75]; $r^2 = 0.026$). In other words, rs6449182 status is not correlated with rs3796863 status [r (Spearman's rho) = -0.004 , $P = 0.97$].

rs6449182

The first section of columns in Table 2 presents the results of analyses for the nine targeted outcomes, showing that rs6449182 was significantly associated with eight of the nine. All associations were in the same direction, indicating that CC individuals had more positive psychological outcomes or behaviors. Specifically, rs6449182 was significantly associated with global evaluation of relationship satisfaction assessed at study entry; after receiving an expression of gratitude, rs6449182 was associated with the perception the expresser was responsive as well as with the specific emotion, loving; after providing an expression of gratitude rs6449182 was associated with the perception that the benefactor was responsive as well as experienced positive emotions (generally) and the specific emotion, loving. rs6449182 was also associated with the behavioral expression of gratitude, either as observed in the videorecorded lab interactions by external judges or as reported by the participant across 14 days: the odds ratio demonstrates that people with the CC genotype were most likely to report thanking the partner for an appreciated event on any given day (see Figure 1 for illustrative data).

rs3796863

The second section of columns in Table 2 presents the results of analyses for the nine targeted outcomes, showing that rs3796863 was significantly associated with one of the nine. Specifically, after providing an expression of gratitude, rs3796863 was significantly associated with the perception that the benefactor was responsive, in the direction indicating that those with the CC genotype were most likely to perceive responsiveness in the benefactor after having expressed gratitude to him or her. In addition, a marginally significant effect emerged for self-reported thanking behavior across 14 days in the same direction, such that those with the CC genotype tended to be most likely to report thanking the partner on a given day. From these rs3796863 analyses, though all coefficients were in the same direction, no other associations exceeded the 0.05 significance threshold.

Table 2 Regression coefficients, significance levels and effect sizes from tests of rs6449182, rs3796863 and putative CD38 expression on the nine targeted outcomes

	CD38					CD38					CD38				
	rs6449182					rs3796863					expression				
	<i>B</i>	95% CI	<i>P</i>	<i>d</i>	<i>B</i>	95% CI	<i>P</i>	<i>d</i>	<i>B</i>	95% CI	<i>P</i>	<i>d</i>			
Relationship satisfaction	-0.15	-0.29, -0.01	0.03	0.39	-0.03	-0.24, 0.17	0.77	0.05	-0.11	-0.23, 0.01	0.07	0.33			
Receiving a gratitude expression															
Perceived responsiveness	-0.24	-0.45, -0.02	0.03	0.40	-0.14	-0.45, 0.18	0.39	0.16	-0.20	-0.37, -0.04	0.02	0.44			
Positive emotions	-0.22	-0.55, 0.11	0.19	0.24	-0.19	-0.66, 0.28	0.43	0.15	-0.20	-0.48, 0.08	0.16	0.26			
Loving	-0.37	-0.69, -0.05	0.02	0.42	-0.09	-0.58, 0.40	0.71	0.07	-0.26	-0.55, 0.02	0.07	0.33			
Expressing gratitude															
Perceived responsiveness	-0.19	-0.35, -0.03	0.02	0.42	-0.37	-0.68, -0.06	0.02	0.43	-0.26	-0.41, -0.12	0.001	0.65			
Positive emotions	-0.31	-0.62, -0.001	0.05	0.36	-0.30	-0.69, 0.09	0.13	0.28	-0.31	-0.54, -0.07	0.01	0.47			
Loving	-0.58	-0.92, -0.24	0.001	0.62	-0.25	-0.62, 0.12	0.18	0.24	-0.46	-0.71, -0.22	0.000	0.69			
Observed praising behavior	-0.21	-0.42, -0.001	0.05	0.39	0.03	-0.20, 0.26	0.79	0.05	-0.12	-0.31, 0.06	0.19	0.26			
Daily expressed gratitude	0.70	0.53, 0.91	0.009	n/a	0.73	0.96, 1.94	0.09	n/a	0.70	0.57, 0.88	0.002	n/a			

B is the unstandardized regression coefficient from multilevel models with *CD38* status as a predictor of the outcome of interest; 95% CI is the 95% confidence interval; *P* is statistical significance level, and the effect size (absolute value) is Cohen's *d*. For daily expressed gratitude, the model accounts for Bernoulli distribution of the outcome (0 or 1), and we therefore present the odds ratio rather than *B*.




Fig. 1 Illustration of linear trend in the percent of days, across 2 weeks, participants reported thanking the partner, by rs6449182 polymorphism. Note that raw data are used in this figure for illustration, but analyses reported in text control for dependence in data across couple-members.

Combined CD38 markers as an index of likelihood of oxytocin influence

The third section of columns in Table 2 presents results of analyses for the nine targeted outcomes, showing that putative CD38 expression was significantly associated with five of the nine (and an additional two approach statistical significance: relationship satisfaction and perception that the partner was loving when receiving an expression of gratitude). All significant (and non-significant) associations were in the same direction, indicating that individuals with presumptive lower CD38 expression had more positive psychological outcomes or behaviors. Specifically, after receiving an expression of gratitude, CD38 expression was significantly associated with the perception that the benefactor was responsive; after providing an expression of gratitude, the CD38 expression variable was associated with the perception that the benefactor was responsive as well as experienced positive emotions (generally) and the specific emotion, loving. Finally, adding ecological validity, from measures across 14 days, lower CD38 expression was associated with a greater likelihood of reporting having thanked the partner for an appreciated event on any given day.²

² Four dependent measures contained outliers (i.e. > 3 s.d. beyond mean). These analyses were re-run with outliers removed, using each index of CD38 expression as predictor; perceived responsiveness after receiving an expression of gratitude became non-significant as predicted by rs6449182 (*P* = 0.11); all other conclusions remained the same.

Alternative explanation

We tested the same pattern of analyses for relational and affective responses to a different conversation topic: disclosure of a personal positive event. Such interactions are positive in valence and responses to them have been associated with global relationship quality (see Gable *et al.*, 2006 for task description and evidence). However, the primary positive emotional theme is joy/enthusiasm rather than gratitude; research demonstrates socially functional distinctions between these emotions (Algoe and Haidt, 2009), and that perceived responsiveness after a partner expressed gratitude better captured the variance in growth in the listener's relationship satisfaction (i.e. promoting bonds) than did perceived responsiveness of the partner after responding to a positive event disclosure (Algoe *et al.*, 2013). No self-report measures after the positive event disclosure interaction (i.e. positive emotions—including loving—or perceived partner responsiveness) was significantly predicted by CD38 status, either when sharing a positive event or when hearing a partner share an event (rs6449181 *P*s = 0.21–0.98, *M P*-value = 0.55; rs3796863 *P*s = 0.06–0.87, *M P*-value = 0.60; CD38 expression *P*s = 0.09–0.98, *M P*-value = 0.60). This suggests that the results for the gratitude expression are not solely due to the positivity of the interaction, and is consistent with theorizing that oxytocin has context-specific effects (Bartz *et al.*, 2011; Feldman, 2012), with the focus of the current investigation on one meaningful context thought to actively promote social bonding.

DISCUSSION

Adult human pair bonds have been posited to be supported by the co-evolution of biological, behavioral and psychological processes (e.g. Hazan and Diamond, 2000; Diamond, 2004; Feldman, 2012). Resting on this assumption, our investigation focused on a putative marker of oxytocin secretion, the behavioral expression of gratitude and psychological responses associated with interactions involving expressions of gratitude, all of which have been closely implicated in promoting the quality of social bonds. In laboratory interactions between people in romantic relationships, and from self-reported behavior in daily life, we found that functional genetic variation in CD38 was associated with the quality and frequency of grateful behavior toward the partner. In addition, this variation also predicted the psychological impact of providing or receiving an expression of gratitude, as well as global relationship satisfaction. Of the two polymorphisms, rs6449182 was associated with a broader spectrum of the building blocks by which gratitude facilitates social bonds. Moreover, when these

uncorrelated markers of oxytocin release were combined into a putative index of *CD38* expression, five of nine targeted effects remained, including thanking behavior in everyday life.

This consistent pattern of findings indirectly linking the oxytocin system with the quality and quantity of expressed gratitude adds to the small but growing body of evidence regarding the behavioral and psychological mechanisms by which oxytocin may influence human pair bonds. Recent theory in relationship science postulates that behaviors associated with prevention of negative outcomes (e.g. fighting well) or alleviation of negative emotions (e.g. receiving social support) are not tapping into the same psychological processes as behaviors associated with the promotion of positive outcomes or emotions (Gable and Reis, 2001). Empirically, negative emotions were very mild in these interactions, whereas positive emotions were relatively high. Consistent with theory, variations in rs6449182 in particular predicted differential positive evaluations of the partner and interaction, whether after providing an expression of gratitude or receiving an expression. Perceived partner responsiveness and positive emotions (e.g. love) after either interaction may have been greater due to consideration of the partner's positive qualities or the partner's actual behaviors in the interaction. Regardless, in so far as *CD38* gene expression affects oxytocin signaling, our results implicate the oxytocin system in the psychological reactions to expressions of gratitude which serve to reward the person for remaining in the relationship. As recent work highlights, it is the acute psychological impact of such interactions that can forecast change in relationship quality (Algoe et al., 2013).

More broadly, our findings are important because the social roles of receiving and providing other-directed positive attention, here enacted through receiving and providing an expression of gratitude, have each recently been linked with consequential downstream outcomes. First, perceiving a partner's responsiveness upon receipt of an expression of gratitude has been associated with downstream relational health benefits (Algoe et al., 2013). The findings regarding the impact of providing an expression of gratitude are interesting in light of the growing evidence for the importance of other-focus for mental health as well as longevity (e.g. Crocker et al., 2009; Konrath et al., 2012). We see our data, examined at the level of a live social interaction, as providing important basic research on potential biological, behavioral and psychological mechanisms for such effects.

Methodologically, integrating live social interactions with measures from daily life, as was done here, represents a valuable path forward for improving understanding of how oxytocin is related to creating social bonds. As reviewed in the introduction, a few studies of structured social interactions in the laboratory generally indicate that oxytocin is associated with beneficial relationship-relevant behaviors. Yet other studies that assess a longer relationship time frame have suggested that oxytocin is associated with distress in the pair-bond relationship (see Taylor et al., 2010). Resolving such controversies will require not only studying the same construct, such as gratitude, across multiple contexts, but also employing different methodologies and manipulations of the oxytocin system to provide convergent validity. In this vein, the current investigation draws attention to the utility of focusing on the gene *CD38*. There is a well-delineated molecular pathway by which rs6449182 can impact cellular signaling, which stands in contrast to commonly studied polymorphisms in the oxytocin receptor gene, such as rs53576, for which there has not been a clearly identified molecular mechanism that could explain how the polymorphism influences signaling. Though the small sample size requires interpretation of these findings as preliminary until replicated, the pattern of associations across a variety of measures that have long been posited to be associated with oxytocin—other-directed positive behavior, intimacy and warm positive feelings (cf. Carter, 1998)—suggests that *CD38* is a

useful marker for future research investigating the role of oxytocin system in the promotion of social bonds.

In addition to being involved in processes other than oxytocin secretion, a limitation of the genetic approach using *CD38* is that because genetic variants are indirect measures of neurotransmitter signaling, it is not yet possible to draw conclusions regarding whether the reported associations with gratitude are due to greater or lesser release of oxytocin in the brain. Mice that do not express *CD38* have low extracellular oxytocin levels (Jin et al., 2007). Similarly, humans with low levels of expression in lymphocytes have lower plasma oxytocin levels (Kiss et al., 2011). In this study, the genotypes were coded based on previously reported effects on *CD38* expression, suggesting that lower *CD38* expression is associated with greater expression of gratitude and greater resulting psychological effects. However, taking the next step to infer that the measures in this study are associated with lower levels of oxytocin release should be made cautiously for several reasons. First, the studies associating rs6449182 and rs3796863 with *CD38* expression (e.g. Jamrozik et al., 2009) were conducted in cells derived from peripheral tissues rather than brain tissue. Yet, it is presumably alterations in *CD38* expression within the brain that are leading to the behavioral and psychological effects seen here. Second, the effects of rs6449182 on gene expression could be influenced by nutritional and cellular factors that were not measured in this study. Third, there are likely to be complex compensatory mechanisms that might counteract the lifelong effects of these polymorphisms. For example, even though mice with deletion of the *CD38* gene have very low levels of oxytocin in the plasma, they actually have very high levels of oxytocin inside the cell because the oxytocin is unable to be released without *CD38* (Jin et al., 2007). Therefore, it will be important in future studies to link these polymorphisms to levels of oxytocin in the cerebrospinal fluid of humans to gain a better understanding of how they might be related to oxytocin signaling and how the aforementioned moderators might impact the effect of rs6449182 on *CD38* expression.

In sum, this research presents the first evidence regarding the biological underpinnings of expressed gratitude in the formation of adult human bonds. Moreover, *CD38* status was associated with behavior and psychological reactions within live interactions, as well as global relationship quality. As an index of oxytocin release, the reliable pattern of results for *CD38* variants across several distinct and theoretically consistent measures provides intriguing evidence for the oxytocin system in solidifying the 'glue' that brings close adult relationship partners closer together.

SUPPLEMENTARY DATA

Supplementary data are available at SCAN online.

REFERENCES

- Algoe, S.B. (2012). Find, remind, and bind: the functions of gratitude in everyday relationships. *Social and Personality Psychology Compass*, 6(6), 455–69.
- Algoe, S.B., Fredrickson, B.L., Gable, S.L. (2013). The social functions of the emotion of gratitude via expression. *Emotion*, 13(4), 605–9.
- Algoe, S.B., Haidt, J. (2009). Witnessing excellence in action: the "other-praising" emotions of elevation, gratitude, and admiration. *Journal of Positive Psychology*, 4(2), 105–27.
- Algoe, S.B., Haidt, J., Gable, S.L. (2008). Beyond reciprocity: gratitude and relationships in everyday life. *Emotion*, 8, 425–9.
- Algoe, S.B., Stanton, A.L. (2012). Gratitude when it is needed most: social functions of gratitude in women with metastatic breast cancer. *Emotion*, 12(1), 163–8.
- Bakermans-Kranenburg, M.J., van IJzendoorn, M.H. (2014). A sociability gene? Meta-analysis of oxytocin receptor genotype effects in humans. *Psychiatric Genetics*, 24(2), 45–51.
- Barrett, J.C., Fry, B., Maller, J., Daly, M.J. (2005). Haploview: analysis and visualization of LD and haplotype maps. *Bioinformatics*, 21, 263–5.
- Bartz, J.A., Zaki, J., Bolger, N., Ochsner, K.N. (2011). Social effects of oxytocin in humans: context and person matter. *Trends in Cognitive Sciences*, 15(7), 301–9.

- Berscheid, E. (1999). The greening of relationship science. *American Psychologist*, 54(4), 260–6.
- Bowlby, J. (1958). The nature of the child's ties to his mother. *International Journal of Psych-Analysis*, 39, 350–73.
- Carter, C.S. (1998). Neuroendocrine perspectives on social attachment and love. *Psychoneuroendocrinology*, 23(8), 779–818.
- Carter, C.S., Porges, S.W. (2013). The biochemistry of love: an oxytocin hypothesis. *EMBO Reports*, 14, 12–6.
- Chapman, J.M., Cooper, J.D., Todd, J.A., Clayton, D.G. (2003). Detecting disease associations due to linkage disequilibrium using haplotype tags: a class of tests and the determinants of statistical power. *Human Heredity*, 56, 18–31.
- Crocker, J., Olivier, M.-A., Nuer, N. (2009). Self-image goals and compassionate goals: costs and benefits. *Self & Identity*, 8, 251–69.
- Diamond, L.M. (2004). Emerging perspectives on distinctions between romantic love and sexual desire. *Current Directions in Psychological Science*, 13(3), 116–9.
- Ditzen, B., Schaer, M., Gabriel, B., Bodenmann, G., Ehlert, U., Heinrichs, M. (2009). Intranasal oxytocin increases positive communication and reduces cortisol levels during couple conflict. *Biological Psychiatry*, 65(9), 728–31.
- Feldman, R. (2012). Oxytocin and social affiliation in humans. *Hormones and Behavior*, 61, 380–91.
- Feldman, R., Gordon, I., Influx, M., Gutbir, T., Ebstein, R.P. (2013). Parental oxytocin and early caregiving jointly shape children's oxytocin response and social reciprocity. *Neuropsychopharmacology*, 38(7), 1154–62.
- Feldman, R., Zagoory-Sharon, O., Weisman, O., et al. (2012). Sensitive parenting is associated with plasma oxytocin and polymorphisms in the OXTR and CD38 genes. *Biological Psychiatry*, 72, 175–81.
- Ferrero, E., Saccucci, F., Malavasi, F. (1999). The human CD38 gene: polymorphism, CpG island, and linkage to the CD157 (BST-1) gene. *Immunogenetics*, 49(7–8), 597–604.
- Fredrickson, B.L. (1998). What good are positive emotions? *Review of General Psychology*, 2, 300–19.
- Gable, S.L., Gonzaga, G.C., Strachman, A. (2006). Will you be there for me when things go right? Supportive responses to positive event disclosures. *Journal of Personality and Social Psychology*, 91, 904–17.
- Gable, S.L., Gosnell, C.L., Maisel, N.C., Strachman, A. (2012). Safely testing the fire alarm: close others' responses to personal positive events. *Journal of Personality and Social Psychology*, 103(6), 963–81.
- Gable, S.L., Reis, H.T. (2001). Appetitive and aversive social interaction. In: Harvey, J., Wenzel, A., editors. *Close Romantic Relationships: Maintenance and Enhancement*. Mahwah, NJ: Lawrence Erlbaum, pp. 169–94.
- Gonzaga, G.C., Turner, R.A., Keltner, D., Campos, B., Altemus, M. (2006). Romantic love and sexual desire in close relationships. *Emotion*, 6(2), 163–79.
- Gouin, J.-P., Carter, C.S., Pournajafi-Nazarloo, H., et al. (2010). Marital behavior, oxytocin, vasopressin, and wound healing. *Psychoneuroendocrinology*, 35, 1082–90.
- Grant, A.M., Gino, F. (2010). A little thanks goes a long way. Explaining why gratitude expression motivate prosocial behavior. *Journal of Personality and Social Psychology*, 98(6), 946–55.
- Greven, K.M., Girdler, S.S., Amico, J., Light, K.C. (2005). Effects of partner support on resting oxytocin, cortisol, norepinephrine, and blood pressure before and after warm partner contact. *Psychosomatic Medicine*, 67(4), 531–8.
- Harlow, H.F. (1958). The nature of love. *American Psychologist*, 13, 673–85.
- Hazan, C., Diamond, L.M. (2000). The place of attachment in human mating. *Review of General Psychology*, 4(2), 186–204.
- Hendrick, S.S. (1988). A generic measure of relationship satisfaction. *Journal of Marriage and the Family*, 50, 93–8.
- Hill, W., Robertson, A. (1968). Linkage disequilibrium in finite populations. *Theoretical and Applied Genetics*, 38(6), 226–31.
- Holt-Lunstad, J., Birmingham, W.A., Light, K.C. (2008). Influence of a "warm touch" support enhancement intervention among married couples on ambulatory blood pressure, oxytocin, alpha amylase, and cortisol. *Psychosomatic Medicine*, 70(9), 976–85.
- Jamrozik, K., Szemraj, Z., Grzybowska-Lzydorzyc, O., et al. (2009). CD38 gene polymorphisms contribute to genetic susceptibility to B-cell chronic lymphocytic leukemia: evidence from two case-control studies in polish caucasians. *Cancer Epidemiology, Biomarkers & Prevention: A Publication of the American Association for Cancer Research, Cosponsored by the American Society of Preventive Oncology*, 18(3), 945–53.
- Jin, D., Liu, H.X., Hirai, H., et al. (2007). CD38 is critical for social behaviour by regulating oxytocin secretion. *Nature*, 446(7131), 41–5.
- Kiss, I., Levy-Gigi, E., Keri, S. (2011). CD 38 expression, attachment style and habituation of arousal in relation to trust-related oxytocin release. *Biological Psychology*, 88(2–3), 223–6.
- Kogan, A., Saslow, L.R., Impett, E.A., Oveis, C., Keltner, D., Saturn, S.R. (2011). Thin-slicing study of the oxytocin receptor (OXTR) gene and the evaluation and expression of the prosocial disposition. *Proceedings of the National Academy of Sciences of the United States of America*, 108(48), 19189–92.
- Kok, B.E., Coffey, K.A., Cohn, M.A., et al. (2013). How positive emotions build physical health: perceived positive social connections account for the upward spiral between positive emotions and vagal tone. *Psychological Science*, 24(7), 1123–32.
- Konrath, S., Fuhrel-Forbis, A., Lou, A., Brown, S. (2012). Motives for volunteering are associated with mortality risk in older adults. *Health Psychology*, 31(1), 87–96.
- Kosfeld, M., Heinrichs, M., Zak, P.J., Fischbacher, U., Fehr, E. (2005). Oxytocin increases trust in humans. *Nature*, 435, 673–6.
- Kubacka, K.E., Finkenauer, C., Rusbult, C.E., Keijsers, L. (2011). Maintaining close relationships: gratitude as a motivator and a detector of maintenance behavior. *Personality and Social Psychology Bulletin*, 37, 1362–75.
- Lambert, N.M., Clark, M.S., Durtschi, J., Fincham, F.D., Graham, S.M. (2010). Benefits of expressing gratitude: expressing gratitude to a partner changes one's view of the relationship. *Psychological Science*, 21, 574–80.
- Lerer, E., Levi, S., Israel, S., et al. (2010). Low CD38 expression in lymphoblastoid cells and haplotypes are both associated with autism in a family-based study. *Autism Research: Official Journal of the International Society for Autism Research*, 3(6), 293–302.
- Lewontin, R.C. (1964). The interaction of selection and linkage. I. General considerations; heterotic models. *Genetics*, 49, 49–67.
- McCullough, M.E., Kilpatrick, S.D., Emmons, R.A., Larson, D.B. (2001). Is gratitude a moral affect? *Psychological Bulletin*, 127(2), 249–66.
- McNulty, J.K., Fincham, F.D. (2012). Beyond positive psychology? Toward a contextual view of psychological processes and well-being. *The American Psychologist*, 67(2), 101–10.
- Munesue, T., Yokoyama, S., Nakamura, K., et al. (2010). Two genetic variants of CD38 in subjects with autism spectrum disorder and controls. *Neuroscience Research*, 67(2), 181–91.
- Polzonetti, V., Carpi, F.M., Micozzi, D., Pucciarelli, S., Vincenzetti, S., Napolioni, V. (2012). Population variability in CD38 activity: correlation with age and significant effect of TNF-alpha -308G>A and CD38 184C>G SNPs. *Molecular Genetics and Metabolism*, 105(3), 502–7.
- Raudenbush, S.W., Bryk, A.S., Cheong, Y.F., Congdon, R.T. (1996). HLM 5. *Hierarchical Linear and Nonlinear Modeling*. Chicago: Scientific Software International.
- Reis, H., Clark, M.S., Holmes, J.G. (2004). Perceived partner responsiveness as an organizing construct in the study of intimacy and closeness. In: Mashek, D.J., Aron, A.P., editors. *Handbook of Closeness and Intimacy*. Mahwah, NJ: Lawrence Erlbaum Associates Publishers, pp. 201–25.
- Riebold, M., Mankuta, D., Lerer, E., et al. (2011). All-trans retinoic acid upregulates reduced CD38 transcription in lymphoblastoid cell lines from autism spectrum disorder. *Molecular Medicine (Cambridge, Mass.)*, 17(7–8), 799–806.
- Saborit-Villarroya, I., Vaisitti, T., Rossi, D., et al. (2011). E2A is a transcriptional regulator of CD38 expression in chronic lymphocytic leukemia. *Leukemia: Official Journal of the Leukemia Society of America, Leukemia Research Fund, U.K.*, 25(3), 479–88.
- Sauer, C., Montag, C., Worner, C., Kirsch, P., Reuter, M. (2012). Effects of a common variant in the CD38 gene on social processing in an oxytocin challenge study: possible links to autism. *Neuropsychopharmacology*, 37(6), 1474–82.
- Schneiderman, I., Zagoory-Sharon, O., Leckman, J.F., Feldman, R. (2012). Oxytocin during the initial stages of romantic attachment: relations to couples' interactive reciprocity. *Psychoneuroendocrinology*, 37, 1277–85.
- Shiota, M.N., Neufeld, S.L., Yeung, W.H., Moser, S.E., Perea, E.F. (2011). Feeling good: autonomic nervous system responding in five positive emotions. *Emotion*, 11(6), 1368–78.
- Smith, T.W., Uchino, B.N., MacKenzie, J., et al. (2013). Effects of couple interactions and relationship quality on plasma oxytocin and cardiovascular reactivity: empirical findings and methodological considerations. *International Journal of Psychophysiology*, 88, 271–81.
- Taylor, S.E., Saphire-Bernstein, S., Seeman, T.E. (2010). Are plasma oxytocin in women and plasma vasopressin in men biomarkers of distressed pair-bond relationships? *Psychological Science*, 21(1), 3–7.
- Trivers, R.L. (1971). The evolution of reciprocal altruism. *Quarterly Review of Biology*, 46, 35–57.
- Uvnas-Moberg, K. (1996). Neuroendocrinology of the mother-child interaction. *Trends in Endocrinology and Metabolism*, 7(4), 126–31.
- Uvnas-Moberg, K. (1998). Oxytocin may mediate the benefits of positive social interaction and emotions. *Psychoneuroendocrinology*, 23(8), 819–35.
- Wigginton, J.E., Cutler, D.J., Abecasis, G.R. (2005). A note on exact tests of Hardy-Weinberg equilibrium. *The American Journal of Human Genetics*, 76(5), 887–93.
- Williams, J.R., Insel, T.R., Harbaugh, C.R., Carter, C.S. (1994). Oxytocin administered centrally facilitates formation of a partner preference in female prairie voles (*Microtus ochrogaster*). *Journal of Neuroendocrinology*, 6, 247–50.